

ALAMEDA COUNTY HEALTH CARE SERVICES AGENCY PUBLIC HEALTH DEPARTMENT

Colleen Chawla, Agency Director **Kimi Watkins-Tartt**, Department Director

Division of Communicable Disease Control and Prevention 1000 Broadway, Ste 500 Oakland, CA 94607 Tel (510) 267-3250 Fax (510) 273-3744 Erica Pan, MD, MPH, Director & Interim Health Officer Sandra Huang, MD, Communicable Disease Controller

Guidance for Alameda County Clinicians: Checklist for Managing Patients Who May Have Novel Coronavirus (2019-nCoV) Infection

Identify patients who may have respiratory illness caused by 2019-nCoV

- Place visible signage requesting visitors with cough and/or fever and recent international travel to immediately notify a hospital staff member.
- Ensure frontline staff including call centers are informed of the current situation with the novel coronavirus and are trained on the facilities triage and communication plan.
- Healthcare providers should screen for travel history on all patients with potentially infectious disease, especially those with fever and acute respiratory illness.

Immediately isolate any persons reporting fever (T >100.4°F or >38°C), lower respiratory symptoms (e.g., cough, shortness of breath), and travel to Wuhan City, China within 14 days prior to illness

- Place surgical mask on patient.
- Place patient in a negative pressure airborne isolation room (If no isolation room available, place patient in private room farthest from areas highly impacted by patients and keep the door closed).

Implement infection control procedures for healthcare workers

- Standard precautions
- Contact precautions (gloves, gown)
- Airborne precautions (N95 mask or PAPR)
- Eye protection (goggles or face shields)

Immediately report the patient to Alameda County Public Health Department

- M-F 8:30-5PM call 510-267-3250
- Afterhours and weekends, call 925-422-7595 and ask to speak to the On-call Public Health Duty Officer
- Public Health will advise on the next steps

Collect specimens for laboratory diagnosis

See instructions below.

Continue medical evaluation and empiric treatment for other causes of respiratory infection or pneumonia as clinically indicated

All patients with suspected 2019-nCoV infection should also be tested for common causes of respiratory infection and pneumonia as clinically indicated. **Testing for other respiratory pathogens should not delay specimen collection for 2019-nCoV testing.**

Discuss plan of care with Public Health prior to discharge

Continue patient isolation and infection control procedures as above.

Novel Coronavirus Specimen Collection Instructions

To increase the likelihood of detecting infection, CDC recommends the collection of specimens from each of these 3 categories: (1) lower respiratory (2) upper respiratory and (3) serum.

1. Upper respiratory tract

- Nasopharyngeal swab <u>AND</u> oropharyngeal swab (NP/OP swab) Use a synthetic fiber swab with plastic shaft. Do not use calcium alginate swabs or swabs with wooden shafts.
 - <u>Nasopharyngeal</u>: insert swab into nostril parallel to the palate, leave swab in place for a few seconds to absorb secretions. Swab both nasopharyngeal areas with the same swab.
 - o <u>Oropharyngeal:</u> swab the posterior pharynx, avoiding the tongue.
 - Place each swab into a separate sterile tube, each with 2-3 ml of viral transport media. Do <u>NOT</u> combine NP/OP swab specimens; keep swabs in separate viral transport media collection tubes. Refrigerate at 2-8°C.

OR

• Nasopharyngeal wash/aspirate or nasal aspirate: collect 2-3 mL into a sterile, leak-proof, screw-cap sputum collection cup or sterile dry container. Refrigerate at 2-8°C.

2. Lower respiratory tract

• **Sputum:** Have the patient rinse the mouth with water and then expectorate deep cough sputum directly into a sterile, leak-proof, screw-cap sputum collection cup or sterile dry container. Refrigerate at 2-8°C.

OR

• **Bronchoalveolar lavage, tracheal aspirate, or pleural fluid:** 2-3 mL in a sterile, leak-proof, screw-cap sputum collection cup or sterile dry container. Refrigerate at 2-8°C.

3. Serum

• Collect 1 tube (5-10 mL) of whole blood in a serum separator tube (minimum 1 mL whole blood is needed for pediatric patients). Refrigerate at 2-8°C.

Note: If specimens cannot be collected at the healthcare facility, do <u>not</u> refer the patient to another facility to obtain specimens. Instead, notify Alameda county Public Health to discuss alternative options.

Specimen Submittal

Alameda County Public Health will provide further instruction on completion of CDC submittal form and coordination of shipping to CDC.

For biosafety reasons, it is <u>not</u> recommended to perform viral isolation in cell culture or initial characterization of viral agents recovered in cultures of specimens from a person under investigation for 2019-nCoV.

See CDC Interim Laboratory Biosafety Guidelines for Handling and Processing Specimens Associated with 2019-nCoV: https://www.cdc.gov/coronavirus/2019-nCoV/lab-biosafety-guidelines.html