

WHAT GAY & BISEXUAL MEN NEED TO KNOW ABOUT MONKEYPOX



Cases have been detected among gay and bisexual men but **not exclusively**

WHAT IS MONKEYPOX?

Monkeypox is a viral infection transmitted through close personal contact, including kissing, sex, and other skin-to-skin contact.

WHAT ARE THE SYMPTOMS?

KEY SYMPTOM



RASH, BUMPS, OR BLISTERS

These may appear anywhere on the body, including the genitals. This may look similar to syphilis, herpes, or other common skin rashes.

OTHER SYMPTOMS



FEVER & HEADACHES



MUSCLE ACHES



SWOLLEN LYMPH NODES

Symptom onset ranges from 5-21 days

WHAT YOU CAN DO



STAY INFORMED

Remain calm. This is a rapidly changing situation. Visit CDC or CDPH websites for up-to-date guidance.



CONTACT

If you have symptoms, call (do not visit) your health care provider, and ask about testing.



TAKE A BREAK

If you have symptoms, stay at home, wear a mask, and cover sores to protect others.

Updated: June 2022



ANYONE CAN GET MONKEYPOX

Blaming any one community may harm public health efforts and cause providers to miss monkeypox in other people.

Get the latest updates & downloadable files from
Gay Sexuality & Social Policy Initiative @ UCLA Luskin
gaysexresearch.com

